DISCLOSURES

This activity is jointly provided by Northwest Portland Area Indian Health Board and Cardea

Cardea Services is approved as a provider of continuing nursing education by Montana Nurses Association, an accredited approver with distinction by the American Nurses Credentialing Center's Commission on Accreditation.

This activity has been planned and implemented in accordance with the accreditation requirements and policies of the Institute for Medical Quality/California Medical Association (IMQ/CMA) through the joint providership of Cardea and Northwest Portland Area Indian Health Board. Cardea is accredited by the IMQ/CMA to provide continuing medical education for physicians.

Cardea designates this live web-based training for a maximum of 1 AMA PRA Category 1 Credit(s)TM. Physicians should claim credit commensurate with the extent of their participation in the activity.





DISCLOSURES

COMPLETING THIS ACTIVITY

Upon successful completion of this activity 1 contact hour will be awarded Successful completion of this continuing education activity includes the following:

- Attending the entire CE activity;
- Completing the online evaluation;
- Submitting an online CE request.

Your certificate will be sent via email If you have any questions about this CE activity, contact Michelle Daugherty at <u>mdaugherty@cardeaservices.org</u> or (206) 447-9538



CONFLICT OF INTEREST

Paulina Deming is on an advisory committee for Gilead. None of the other planners or presenters of this CE activity have any relevant financial relationships with any commercial entities pertaining to this activity.



Acknowledgement

This presentation is funded in part by:

The Indian Health Service HIV Program and The Secretary's Minority AIDS Initiative Fund





HCV and Children:

Part 1 - What do we know?

Jacqueline L. Fridge, MB ChB, MBA Division Chief Pediatric Gastroenterology, Hepatology and Nutrition, University of New Mexico *jfridge@salud.unm.edu*

01/17/2018

Presentation prepared by: Date prepared: 5

Ø

Objectives

- Discuss prevalence of HCV in children
- Discuss screening children and youth for HCV



Prevalence

Worldwide 170,000,000 million infected: 5 million children

- Highest prevalence in Egypt; 3% prevalence in children
- US:
 - 4 million adults living with HCV infection; 1-2% adults infected
 - M>F; rising incidence in baby boomers 50-69 y/o
 - Non-Hispanic black 2 x more likely & Native Americans 3 x more likely to be diagnosed as non-Hispanic whites.
 - 0.1-0.2% children infected:
 - 0.15% of 6-11 year olds
 - 0.4% of 12-19 year olds
 - 23,000 to 46,000 children in US with HCV

Lee 2015 WMJ; Rayne-Greenow 2015 J Peds & Child Health



What about New Mexico?

Hepatitis C Rates in New Mexico by County, 2015



Grouping: Equal Groups (Quantile)

Dates are from NM IBIS on 5/9/2016

Drawing population data from NM 2016 Census:

- Estimate 982 to 1,965 children < age 18 have been exposed to HCV
- Estimate 246to 1,768 children are chronically infected



Vertical transmission

Prevalence of HCV infection in pregnant women (HCV Ab or pcr +ve) 0.75% (range 0.5-1.7%) Overall risk of transmission 5%:

- 1.7% per pregnancy if mom HCV Ab positive
- 4.3% per pregnancy if mom HCV PCR positive
- 7.1% per pregnancy if mom HCV PCR positive at least twice during pregnancy or around the time of delivery
- 2-3 fold higher if mom co-infected with HIV
- 36% if maternal viral load ≥ 1,000,000 iu/ml

Unknown at what stage of pregnancy transmission occurs

Thought that method of transmission differs from that of HIV

All genotypes can cause MTCT (mother to child transmission), but limited quaisispecies are noted

As HCV is present in monocytes too, suggestion HLA similarity between mother and child may facilitate persistence of maternal cells in newborn blood

Accounts for 7,500 new cases HCV in US per year

HCV transmission from an infected father via sperm has been demonstrated

- (Ma et al. 2016 Clin Lab)

HCV can infect placental cells



Vertical transmission risk factors

Maternal risk factors:	Infant risk factors:
Coinfection with HIV	Fetal scalp vein monitoring?
IV drug use?	Fetal anoxia at the time of delivery (cord blood pH)
Prolonged rupture of membranes > 6 hours	Protective HLA types (mismatch favorable?)
High maternal HCV load, >600,0000 IU/ml, but genotype not a factor	Higher risk HLA types: HLA-DRB1*10
Risk of amniocentesis not known (HCV detected in amniotic fluid)	Higher risk for females vs males?
No reduction of risk with elective LSCS (Cochrane)	
Avoid large vaginal tears with vaginal delivery (episiotomy not a risk factor?)	
Maternal HLA: HLA-DRB104 protective	
Not related: maternal age; parity; cigarette smoking; alcohol use	
No benefit to withholding breast feeding unless nipples are bleeding, active mastitis (CDC) or mom has post partum HCV flare with jaundice	

TORCH infections

Infection	US prevalence of congenitally acquired disease
Toxoplasmosis	10-33/100,000 live births
Treponema pallidum	7.8/100,000 live births
CMV	800/1,000,000 live births
Hepatitis B	<0.1/1,000,000 live births
Hepatitis C	<0.1/100,000 live births
HIV	162 infants/yr, 2010

Neu et al, 2015, Clin Perinatol



HCV Screening of Pregnant women

Studies show not cost effective to screen all pregnant women for HCV

Screen high risk pregnant women

Screening transaminases is not helpful as they often improve in 2nd & 3rd trimester

ACOG recommendations

- Screen pregnant women with risk factors such as:
 - IVDU (but self-reporting is unreliable)
 - HIV

Also:

- HBV
- Blood transfusions
- Clotting factors

10 year study screening all pregnant women at one center in UK:

- 35,000 women screened
- 136 positive (0.38%) HCV Ab positive
- 78 active viremia (0.22%)
- 44 new diagnosis (0.12%)
 - 3 vertical MTCT in this group (6.8%)
 - 19 treated post partum (IFN & ribavirin) with 74% SVR
- Cost effective in their algorithm
- (Selvapatt et al 2015 J of hepatology)

FDA approval for use of sofosbuvir & ledipasvir in pregnancy may be coming soon.



Proposal to screen all pregnant women and offer treatment with DAAs before next pregnancy? (2016 Aebi-Popp et al. J Virus Eradication)

Pregnant women with HCV infection have worse maternal and fetal outcomes:

- Gestational diabetes
- Preeclampsia
- Growth restriction
- Antepartum hemorrhage
- Preterm birth
- Gestational cholestasis
- Physiologic state of later pregnancy may favor viral replication

Infants of HCV positive mother have worse outcomes independent of IVDU:

- Low birth weight
- Ventilator assistance
- NICU admission
- Congenital anomalies?



Other methods of infection of children

- IV drug use /needle sharing
- High-risk sexual behaviors (negligible risk for sexual intercourse in stable relationship)
- Household exposure <2%
- Tattooing or body piercing





Questions for ECHO Providers

- Why are so few children referred to Peds GI clinic for HCV?
 - Are they diagnosed and not referred?
 - Are they not diagnosed?
 - Other reason?
- When treating an adult patient with HCV do you have a protocol for screening that person's children for possible HCV infection?
 - Yes
 - No
 - Not a formal one but I think about it!
- What are the barriers to screening children and youth?
 - Time?
 - I'm not their PCP
 - Parents resistant
 - Other?



Whom to test? Identifying at risk & infected children

Group	Screening
Adolescents who have injected illicit drugs in the recent and remote past, including those who injected only once and do not consider themselves to be drug users	Antibody
 Children and adolescents with conditions associated with a high prevalence of HCV infection including: Persons with human immunodeficiency virus (HIV) infection Persons who have ever been on hemodialysis 	Antibody or RNA
Children born to HCV-infected mothers	Antibody >18 months RNA ≤ 18 months
Present sexual partners of HCV-infected persons	Antibody
Children or adolescents with chronically elevated transaminases	Antibody
 Children and adolescents with signs or symptoms of hepatitis: Jaundice, dark urine, pale stool, fatigue, nausea and vomiting, abdominal pain, especially in the RUQ, loss of appetite, low-grade fever, joint pain 	Antibody and/or RNA
 Children and adolescents with intrafamilial exposure: Siblings, parents, household contacts 	Antibody

Whom to test?

Group	Screening	
 Children and adolescents emigrating from a region with high prevalence of HCV infection: North Africa/Middle East: (Egypt, Iraq, Yemen) Central Sub-Saharan Africa: (Burundi, Democratic Republic of the Congo, Gabon) Central Asia: (Georgia, Mongolia, Uzbekistan) Eastern Europe: (Moldova, Russia, Ukraine)[3] 	Antibody	
Adolescents incarcerated in a correctional institution	Antibody	
Adolescents who received tattooing and body piercing in unregulated settings	Antibody	
Children or adolescents after a needle stick injury or mucosal exposure to HCV-positive blood.	Antibody	
Adolescents diagnosed with a new sexually transmitted disease **	Antibody	



Whom to Test? References/notes

Note: All of the positive anti-HCV antibody tests should be followed up with an HCV RNA test to determine active infection.

The serum HCV RNA can be tested before 18 months of age; however, infants should be at least 2 months old.[4] If serum HCV RNA is positive in early infancy, it should be rechecked after 12 months of age to determine presence of chronic hepatitis C (CHC).

1. Squires, J.E. and W.F. Balistreri, *Hepatitis C virus infection in children and adolescents*. Hepatology Communications, 2017. **1**(2): p. 87-98.

2. Mack, C.L., et al., NASPGHAN practice guidelines: Diagnosis and management of hepatitis C infection in infants, children, and adolescents. J Pediatr Gastroenterol Nutr, 2012. **54**(6): p. 838-55.

3. Petruzziello, A., et al., *Global epidemiology of hepatitis C virus infection: An up-date of the distribution and circulation of hepatitis C virus genotypes.* World J Gastroenterol, 2016. **22**(34): p. 7824-40.

4. Polywka, S., et al., Accuracy of HCV-RNA PCR tests for diagnosis or exclusion of vertically acquired HCV infection. J Med Virol, 2006. **78**(2): p. 305-10.

** need ref for this



Natural History in Children (vertical transmission)

- 1) Spontaneous clearance: 25-40% perinatally infected
 - Clearance of transient neonatal viremia vs. true AHCV with increased transaminases?
 - Typically by 2 years of age, can occur as last as 7 y/o.
 - (adults only clear within the first 6 months of infection)
 - Greatest clearance genotype 3 and IL-28B polymorphisms (response of interferon to viral antigens) and other genotype polymorphisms
 - Positive association between increased ALT in first 2 years and viral clearance

2) Chronically infected: no clearance by 2 y/o

- A) Asymptomatic 50%:
 - normal transaminases
 - intermittent viremia
- B) Chronic active infection 30%:
 - Frequently abnormal transaminases
 - Hepatomegaly
 - Persistent viremia
 - Bridging fibrosis, cirrhosis O.5 to 10% (1-2%)
 - hepatocellular carcinoma can occur
 - Risk factors for fibrosis include positive anti LKM Ab
- 80% have minimal to no liver fibrosis by 18 y/o
- ALT level is a poor predictor of liver fibrosis
- Very rare reason for liver transplantation children



Natural history in children (non-vertical infection)

- Similar to adults
- 6-12% spontaneous clearance, but some studies show higher rates; highest for Genotype 3.
- 80% develop chronic HCV
 - 1-2% develop cirrhosis in childhood
 - Chronic liver disease with cirrhosis in 20-30 years
 - Risk factors for more severe disease:
 - Obesity
 - Treatment for childhood cancer
 - Congenital anemia requiring chronic transfusion
 - Coinfection with HBV or HIV
 - Social factors:
 - Homelessness
 - IVDU & alcohol use
 - incarceration

- HCV associated comorbidities (vertical and non-vertical HCV):
 - Glomerulonephritis membranoproliferative
 - No reports of cryoglobulinemia or lymphoma
 - (cryoglobulins and autoantibodies can be detected but generally not clinically significant)
 - CNS infection? -> learning impairment
 - Decreased QOL



Criteria for diagnosis?

Criteria to rule out infection?

Positive confirmed HCV antibody at least once in child \geq 18 months

• Confirm with positive HCV RNA PCR

Positive HCV RNA test, PCR at any age > 3 months old (2 months)

- High rate of temporary false positives in children < 3 months
- Peripheral transient viremia typically occurs days 0-5 of life
- Cord blood viremia not indicative of neonatal infection
- Should be rechecked after 12 months of age to confirm CHCV

Positive HCV RNA at 2-6 months and positive RNA and Ab at 18-24 months

Criteria to rule out vertical transmission:

 Two negative HCV-PCR tests at least 2-3 months apart

Criteria for HCV clearance after chronic infection:

 Two negative HCV-PCR tests at least 6 months apart



Evaluation

- HCV PCR viral load quantitative
- HCV genotype
- Transaminases; GGT;
- Liver function: albumin; bilirubin; PT & inr; (+/- glucose; fibrinogen; ammonia)
- BUN & Cr (GFR) +/- urinalysis
- Nutrition:
 - Fat soluble vitamin levels: A, D, E and PT & inr
 - CBC
- Thyroid: TSH, FT4
- Baseline abdominal USS

Evaluate for other liver disease?

- Infection:
 - HIV antibody
 - Hep B sAb; aAg; eAg; cAb
 - TORCH
 - CMV & EBV
 - Adeno
- Infants:
 - Genetic:
 - Alpha-1-antitrypsin deficiency
 - Cystic fibrosis
 - Inborn errors of metabolism
- Teens:
 - NAFLD/steatohepatitis
 - Autoimmune hepatitis
 - Gilbert's syndrome
 - Wilson's Disease

When to treat

- Medical indications:
 - Children with aggressive liver disease (with fibrosis)
 - Persistently elevated transaminases
 - Children with genotype 2 or 3
- Treat all children?
 - 26 fold increased risk of liver related death if HCV acquired in childhood
 - Economic burden of years of decades of infection
 - Prevent transmission
- Family treatment?
- Goals of treatment:
 - Eradicate infection
 - Preventing end stage liver disease and HCC
 - Removing stigma of HCV infection
 - Decrease global burden of disease/ HCV eradication

- Exclusions:
 - No treatment < 3 years of age
 - Severe co-morbid disease
 - Active IVDU or alcohol use? (Patient-Centered Models of HCV Care for People Who Inject Drugs (HERO))
 - Active psychiatric disease
 - Seizures
 - Incarceration?
 - Solid organ transplant?





Treatment:

Currently approved for children under 12 y/o

- Pegylated interferon and ribavirin (2008):
 - PEG-IFN-α sq q week and ribavirin PO BID
 - Genotype 1 or 4 45% achieve sustained viral response (SVR = no virus detected in blood by HCV-PCR) at 24 weeks after completing treatment) after 48 weeks of therapy
 - Genotype 2 or 3 SVR 80% after **24 weeks** of therapy
 - HIV or HBV coinfection require 48 week treatment irrespective of genotype
- Better SVR:
 - Genotype 2 & 3
 - Genotype 1 with viral load <600,000 IU/ml or <2 x 10⁶ copies/ml
 - Two positive genetic polymorphisms of IL-28B receptor
- Worse SVR with obesity
- IFN-α: cytokine which activates the innate antiviral immune system
- Ribavirin: oral nucleoside analogue with antiviral and immunomodulatory effects



April 2017: DAA approved for children 12-17 y/o

Harvoni (ledipasvir and sofosbuvir) 90mg/400 mg tablet:

- Pediatric patients 12 years of age and older or weighing at least 77 pounds (35 kilograms)
- HCV genotype 1, 4, 5 or 6 infection without cirrhosis or with mild cirrhosis.
 - Genotype 1, Tx-naïve patients: 1 tab PO qd x 12 wk
 - Genotype 4, 5 or 6, Tx-naïve patients: 1 tab PO qd x 12 wk

Sovaldi (sofosbuvir) 400 mg tablet:

- Sovaldi in combination with ribavirin
- Pediatric patients 12 years of age and older or weighing at least 77 pounds (35 kilograms)
- HCV genotype 2 or 3 HCV infection without cirrhosis or with mild cirrhosis.
 - Genotype 2: 400 mg PO qd w/ribavirin x 12 wk
 - Genotype 3: 400 mg PO qd w/ribavirin x 24 wk
 - Ribavirin 200 mg capsule or tablet (do not open/crush/chew cap):
 - 35-46kg: 15mg/kg/day PO divided BID
 - 47-49kg: 600 mg/day PO divided BID
 - 50-65kg: 800 mg/day PO divided BID
 - 66-80kg: 1,000 mg/day PO divided BID
 - > 80kg: 1,200 mg/day PO divided BID
- Harvoni and Sovaldi: Black Box warning for Hepatitis B reactivation
- Ribavirin: Black Box warning for hemolytic anemia and teratogenic/embryocidal effects



Questions for ECHO providers

- Would you feel comfortable treating children with DAAs?
- What support would you need?
- Should all children with HCV be referred to Peds GI clinic?



Project HepCCY – UNM IRB approved registry

Hepatitis C in Children and Youth of NM







Project ECHO HCV Collaborative

End of Presentation

Questions?













Evaluation and Certificates

• Please use the link below to complete the learner evaluation. This link will also be emailed to you within a few days. Please check your junk and spam email folders if you don't receive it.

http://sgiz.mobi/s3/July-GP

